

PCT

WORLD INTELLECTUAL PROPERTY
ORGANIZATION
International

INTERNATIONAL APPLICATION PUBLISHED UNDER

WO 9605739A1

(51) International Patent Classification ⁶ : A23K 1/165, 1/18	A1	(11) International Publication Number: WO 96/05739 (43) International Publication Date: 29 February 1996 (29.02.96)
(21) International Application Number: PCT/EP95/03277 (22) International Filing Date: 17 August 1995 (17.08.95) (30) Priority Data: 9416841.6 19 August 1994 (19.08.94) GB (71) Applicants: FINNFEEDS INTERNATIONAL LIMITED [GB/GB]; Market House, Ailesbury Court, High Street, Marlborough, Wiltshire SN8 1AA (GB). GENENCOR INTERNATIONAL INC. [US/US]; 4 Cambridge Place, 1870 Winton Road South, Rochester, NY 14618 (US). (72) Inventors: BEDFORD, Michael, Richard; Finnfeeds Interna- tional Ltd., Market House, Ailesbury Court, High Street, Marlborough, Wiltshire SN8 1AA (GB). MORGAN, An- drew, John; Finnfeeds International Ltd., Market House, Ailesbury Court, High Street, Marlborough, Wiltshire SN8 1AA (GB). CLARKSON, Kathleen; Genencor International Ltd., 180 Kimball Way, South San Francisco, CA 94080 (US). SCHULZE, Hagen, Klaus; Finnfeeds International Ltd., Market House, Ailesbury Court, High Street, Marlbor- ough, Wiltshire SN8 1AA (GB). (74) Agents: MITCHELL, Alan et al.; Hoffmann, Eitle & Partner, Arabellastrasse 4, D-81925 München (DE).	(81) Designated States: AU, CA, CN, FI, JP, NO, NZ. Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments</i>	
(54) Title: AN ENZYME FEED ADDITIVE AND ANIMAL FEED INCLUDING IT		
(57) Abstract		
<p>An enzyme feed additive is provided comprising a xylanase, a protease, and optionally a β-glucanase. The ratio of the units of xylanase activity per unit amount of the feed additive to the units of β-glucanase activity per same unit amount of the feed additive is 1:0-0.25. Preferably, the xylanase is the low pI xylanase and/or the high pI xylanase obtained from <i>Trichoderma longibrachiatum</i>. Preferably, the protease is a mutant subtilisin comprising a substitution at the amino acid residue position equivalent to tyr+217 of <i>St</i>(<i>Bacillus amyloliquefaciens</i>, subtilisin with leucine.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

AN ENZYME FEED ADDITIVE AND ANIMAL FEED INCLUDING IT

The present invention relates to an enzyme feed additive, and in particular to such an additive which can decrease the feed conversion ratio of a cereal-based animal feed.

Improvements in animal feeds to enable animals to digest them more efficiently are constantly being sought. One of the main concerns is to improve the feed conversion ratio (FCR) of a feed without increasing its cost per unit weight. The FCR is the ratio of the amount of feed consumed relative to the weight gain of an animal. A low FCR indicates that a given amount of feed results in a growing animal gaining proportionately more weight. This means that the animal is able to utilise the feed more efficiently. One way in which the FCR of a feed can be improved is to increase its digestibility.

There are various constraints on the digestibility of the nutritional components of a feed such as its starch, fat, protein and amino acid contents. These constraints include:

- (i) the viscosity of materials present in the animal's gut. Such viscosity is due, at least in part, to soluble non-starch polysaccharides such as mixed-linked β -glucans and arabinoxylans;

in cereals. Such entrapment is caused by the high levels of non-starch polysaccharides in the cell walls of cereals which are relatively resistant to break-down by the animal's digestive system. This prevents the nutrients entrapped within the cells from being nutritionally available to the animal; and

(iii) a deficiency in endogenous enzyme activity, both of the animal and of the gut microbial population particularly in a young animal.

The above problems which interfere with digestibility are particularly noticeable in the case of cereal-based diets, such as those having a high wheat content.

Due to the problem of poor digestibility of nutrients from the feed, it is normally necessary to formulate feeds to contain significant amounts of energy providing materials in order to meet the nutritional demands of animals. Such energy providing materials conventionally include starch, fat, sugars, fibre etc. The requirement of including these energy providing materials, or sources of such materials, in a feed adds a considerable extra cost which is disadvantageous from an economic view point.

In an attempt to solve the problem of poor digestibility of cereal-based feeds, it is known to include enzyme supplements such as β -glucanases or xylanases in animal feeds. For example, WO 91/04673 discloses a feed additive for

alleviating malabsorption syndrome in poultry which causes reduced digestion. The additive includes a cellulase and a xylanase. JP-A-60-75238 discloses a feed for domestic animals which contains an enzyme cocktail including protease-, cellulase-, amylase- and lipase-activities. This reference speculates that these various enzyme activities enable fermentation microbes to grow and these become useful nutritional components of the feed.

Cellulases (i.e. the cellulase system) are enzyme compositions which hydrolyze cellulose (β -1,4-D-glucan linkages) and/or derivatives thereof (e.g. phosphoric acid swollen cellulose) and give as primary products glucose, cellobiose, cellooligosaccharides, and the like. A cellulase system produced by a given microorganism is comprised of several different enzyme classifications including those identified as exo-cellobiohydrolases (EC 3.2.1.91) ("CBH"), endoglucanases (EC 3.2.1.4) ("EG"), and β -glucosidases (EC 3.2.1.21) ("BG") (Schulein, M., 1988). Moreover, these classifications can be further separated into individual components. For example multiple CBH components and EG components have been isolated from a variety of bacterial and fungal sources such as the cellulase enzyme complex of *Trichoderma longibrachiatum* which consists of two exo-cellobiohydrolases, CBHI and CBHII, at least three endoglucanases, EGI, EGII and EGIII, and at least one β -glucosidase. The endoglucanases and exo-cellobiohydrolases are considered to act synergistically in the hydrolysis of cellulose to small soluble sugars.

the action of β -glucosidase. In addition to hydrolyzing the β -1,4 linkages in cellulose, endo-1,4- β -glucanase (EC 3.2.1.4) will also hydrolyze 1,4 linkages in β -glucans also containing 1,3-linkages. The endoglucanases act on internal linkages to produce cellobiose, glucose and cello-oligosaccharides. The exo-cellobiohydrolases act on non-reducing ends of cellulose polymers to produce cellobiose as the principal product.

Organisms which produce or express cellulase enzyme complexes often also express xylanase activity. For example, two different xylanase enzymes have been identified which are produced by *T. longibrachiatum*. The purification of these two different xylanases, one referred to as high pI xylanase (having a pI of about 9.0) and the other referred to as low pI xylanase (having a pI of about 5.2), as well as the cloning and sequencing of the gene for each xylanase is described in detail in WO 92/06209. Figure 16 of this document sets out the deduced amino acid sequences for both the low pI and high pI gene products. Example 22 also teaches how to create *T. longibrachiatum* strains which over-express the low pI and high pI xylanase genes and which are unable to produce some or all of the non-xylanase cellulase components normally associated with *T. longibrachiatum* such as CBHI, CBHII, EGI, EGII, EGIII and BG.

As mentioned above, the use of cellulases and xylanases in animal feeds is known as is the use of proteases. It is found however that many naturally occurring sources of xylanase contain a significantly greater relative activity of

β -glucanase. For example in natural strains of *T. longibrachiatum*, the ratio of xylanase activity to β -glucanase activity is of the order 1:5. Most surprisingly, it has been found that when a xylanase is included in a cereal-based diet at or around its optimum dosage level, the co-presence of enzymes possessing β -glucanase activity increases the FCR of the feed which is of course disadvantageous. In view of this surprising finding, it is concluded that enzymes having β -glucanase activity are not only unnecessary in feeds supplemented with a xylanase, but their presence is in fact detrimental to the benefits obtained from the presence of the xylanase and/or protease.

In the description and claims which follow, reference is made to units of xylanase activity, units of protease activity, and units of β -glucanase activity. These three activities as used in the present specification are measured by the following three assay methods.

Assay Method for Xylanase Activity

One unit of xylanase activity is the amount of enzyme which liberates one μ mol of reducing sugars (expressed as xylose equivalents) from the substrate in one minute under the conditions described.

Reagents

1. 1% (w/v) xylan substrate

Add 10 ml of 0.5 M sodium hydroxide to 1.0 g of xylan (Fluka 95590). Mix for 30 minutes with a magnetic stirrer. Add about 40 ml of 0.05 M sodium acetate buffer, pH 5.3. Adjust pH to 5.3 with 1 M acetic acid. Fill to 100 ml with 0.05 M sodium acetate buffer, pH 5.3. Substrate should be mixed all the time when used.

2. 1 M acetic acid

Pipette 5.7 ml of glacial acetic acid into a volumetric flask and fill to 100 ml with distilled water.

3. 0.05 M sodium acetate buffer, pH 5.3

A. Dissolve 4.1 g of sodium acetate in distilled water and fill to 1000 ml with distilled water.

B. Dissolve 3.0 g of glacial acetic acid in distilled water and fill to 1000 ml with distilled water.

Adjust the pH of solution A to pH 5.3 with solution B.

4. Dinitrosalicylic acid (DNS) reagent

Suspend 20.0 g of 3,5-dinitrosalicylic acid in about 800 ml of distilled water. Add gradually 300 ml of sodium hydroxide solution (32.0 g NaOH in 300 ml of distilled water) while stirring continuously. Warm the suspension in a water bath (the temperature may not exceed +48°C) while stirring until the solution is clear. Add gradually 600 g of potassium sodium tartrate. Warm the solution (the temperature may not exceed +48°C) if needed until the solution is clear.

Fill to 2000 ml with distilled water and filter through a coarse sintered glass filter.

Store in a dark bottle at room temperature. The Reagent is stable for a maximum of 6 months.

Procedure

1. Enzyme sample

1 ml of enzyme dilution (in 0.05 M sodium acetate buffer, pH 5.3) is equilibrated at +50°C. Add 1 ml of xylan substrate, stir and incubate at +50°C for exactly 30 minutes.

5 minutes. Cool the reaction mixture in a cold water bath to room temperature and measure the absorbance at 540 nm against distilled water.

2. Enzyme blank

Incubate 1 ml of xylan substrate at +50°C for 30 minutes. Add 3 ml of DNS-solution and stir. Add 1 ml of enzyme dilution (in 0.05 M sodium acetate buffer, pH 5.3) and stir. Boil the mixture for exactly 5 minutes. Cool the reaction mixture in a cold water bath to room temperature and measure the absorbance at 540 nm against distilled water.

The absorbance difference between the enzyme sample and enzyme blank should be 0.3-0.5.

3. Standard curve

Prepare standard solutions from anhydrous xylose in 0.05 M sodium acetate buffer, pH 5.3. Xylose concentration in the standards should be 0.05-0.5 mg/ml. Pipette 1 ml of standard solution, 1 ml of xylan substrate and 3 ml of DNS-reagent into a test tube. Stir and boil for exactly 5 minutes. Cool in a cold water bath to room temperature and measure the absorbance at 540 nm against

standard blank. In the standard blank, xylose solution is replaced by 1 ml of 0.05 M sodium acetate buffer, pH 5.3. Otherwise standard blank is treated like xylose standard.

Plot xylose concentration as a function of absorbance. New standard curve is prepared for every new DNS-reagent.

Calculation

The xylanase activity of the sample is calculated according to the following equation:

$$\text{Activity (U/g)} = \frac{([A(X) - A(O)] \times k + C) \times 1000 \times Df}{MW_{\text{xy1}} \times t}$$

wherein:

A(X)	=	absorbance of the enzyme sample
A(O)	=	absorbance of the enzyme blank
k	=	the slope of the standard curve
C.	=	the intercept of xylose standard curve
1000	=	factor, mmol -> µmol
Df	=	dilution factor (ml/g)
MW _{xy1}	=	molecular weight of xylose (150.13 mg/mmol)
t	=	reaction time (30 minutes)

Assay Method for Protease Activity

One unit of protease activity is the amount of enzyme which liberates from the substrate one microgram of phenolic compound (expressed as tyrosine equivalents) in one minute under the conditions described.

Reagents 1. 0.6% (w/v) casein substrate

Weigh 0.6 g of dry Hammarsten Casein (Merck 2242) into a 200 ml beaker. Moisten with a small amount (about 5 ml) of distilled water. When casein is thoroughly moistened add 20 ml of 0.2 M disodium hydrogen phosphate solution. Warm the mixture at +60°C with stirring until casein dissolves and an opal solution is obtained. Add 60 ml of distilled water and if needed 1-2 drops of octyl alcohol (anti-foam agent; similar products can be used). After cooling to room temperature, adjust the pH to 7.5 with 0.5 M sodium hydroxide and 1 M lactic acid. Transfer solution into a volumetric flask and fill to 100 ml with distilled water.

Substrate solution is usable for one week if stored in a cold room.

2. 0.2 M Na_2HPO_4 solution

Dissolve 17.80 g of disodium hydrogen phosphate dihydrate in distilled water and fill to 500 ml with distilled water.

3. 0.02 M NaCl solution

Dissolve 1.168 g of sodium chloride in distilled water and fill to 1000 ml with distilled water.

4. Precipitation reagent (TCA)

Dissolve 18.80 g of trichloroacetic acid (CCl_3COOH), 18.10 g of anhydrous sodium acetate (CH_3COONa) and 18.80 g of acetic acid (CH_3COOH) in distilled water and fill to 1000 ml with distilled water.

5. Phenol reagent

Mix one (1) part of Folin-Ciocalteu phenol reagent with one (1) part of distilled water just prior to the assay.

6. 0.55 M Na_2CO_3 solution

distilled water and fill to 1000 ml with distilled water.

Procedure

1. Enzyme sample

Equilibrate 1 ml of enzyme dilution (in 0.02 M NaCl solution) at + 40°C (for about 5 minutes). Add 5 ml of equilibrated casein substrate, stir and incubate at + 40°C for exactly 30 minutes. Add 5 ml of precipitation reagent and stir. Incubate at + 40°C for exactly 30 minutes and filter immediately with filter paper (Whatman 1 or Macherey Nagel 640 we).

Pipette 2 ml of filtrate, 5 ml of 0.55 M Na_2CO_3 solution and 1 ml of phenol reagent. Stir and incubate at +40°C for 30 minutes. Cool to room temperature and measure the absorbance at 660 nm against distilled water.

2. Enzyme blank

Equilibrate 1 ml of enzyme dilution (in 0.02 M NaCl solution) at +40°C (for about 5 minutes). Add 5 ml of precipitation reagent, stir and incubate at +40°C for exactly 30 minutes. Add 5 ml of casein substrate, stir and incubate at +40°C for exactly 30 minutes.

Filter immediately with filter paper (Whatman 1 or Macherey Nagel 640 we).

Treat the filtrate as the enzyme sample.

The absorbance difference between the enzyme sample and the enzyme blank should be 0.2-0.5.

3. Standard curve

Prepare a tyrosine stock solution by weighing 10 mg of L-tyrosine into a volumetric flask, dissolve in 0.02 M NaCl solution and fill to 100 ml with 0.02 M NaCl solution.

Prepare dilutions from tyrosine stock solution in 0.02 M NaCl solution as follows:

1:50	=	2 µg/ml
1:20	=	5 µg/ml
1:10	=	10 µg/ml
1:5	=	20 µg/ml
1:3	=	33 µg/ml
1:2	=	50 µg/ml

Pipette 2 ml of each tyrosine dilution, 5 ml of 0.55 M Na₂CO₃ solution and 1 ml of phenol

measure the absorbance at 660 nm against distilled water.

Plot tyrosine concentration as a function of absorbance.

Calculation The protease activity of the sample is calculated according to the following equation:

$$\text{Activity (U/g)} = \frac{[A(X) - A(O)] \times k \times F \times Df}{t}$$

wherein:

A(X)	=	absorbance of the enzyme sample
A(O)	=	absorbance of the enzyme blank
k	=	the slope of the standard curve
F	=	reaction dilution factor (= 11)
Df	=	dilution factor (ml/g)
t	=	reaction time (30 minutes)

Assay Method for β -Glucanase Activity

One unit of β -glucanase activity is the amount of enzyme which liberates one μmol of reducing sugars (expressed as glucose equivalents) from the substrate in one minute under the conditions described.

Reagents

1. 1.0% (w/v) β -glucan substrate

Moisten 1.0 g of mixed-linked β -(1,3)(1,4)-glucan (Biocon Biochemicals Ltd.) with 10 ml of ethanol. Add about 80 ml of distilled water and warm up to boil. Continue boiling with vigorous stirring until β -glucan is dissolved and a turbid solution is obtained. Cool the turbid solution to room temperature continuously stirring and adjust the β -glucan concentration to 1.0% (w/w) by adding distilled water. Filter through a glass fibre filter paper.

The substrate can be used immediately. The substrate is usable for two days if stored in a cold room.

2. 0.1 M sodium acetate buffer, pH 5.0

A. Dissolve 8.2 g of anhydrous sodium acetate in distilled water and fill to 1000 ml with distilled water.

B. Dissolve 6.0 g of glacial acetic acid in distilled water and fill to 1000 ml with distilled water.

Adjust the pH of solution A to 5.0 with solution B.

3. Dinitrosalicylic acid (DNS) reagent

Suspend 20.0 g of 3,5-dinitrosalicylic acid in about 800 ml of distilled water. Add gradually 300 ml of sodium hydroxide solution (32.0 g of NaOH in 300 ml of distilled water) while stirring continuously. Warm the suspension in a water bath (the temperature may not exceed +48°C) while stirring until the solution is clear. Add gradually 600 g of potassium sodium tartrate. Warm the solution (the temperature may not exceed +48°C) if needed until solution is clear.

Fill to 2000 ml with distilled water and filter through a coarse sintered glass filter.

Store in a dark bottle at room temperature. The reagent is stable for a maximum of 6 months.

Procedure

1. Enzyme sample

Equilibrate 1 ml of enzyme dilution (in 0.1 M sodium acetate buffer, pH 5.0) at -30°C. Add 1 ml of β -glucan substrate, stir and incubate

SUBSTRATE: 0.1 M β -glucan

at +30°C for exactly 10 minutes. Add 3 ml of DNS-reagent, stir and boil the reaction mixture for exactly 5 minutes. Cool the reaction mixture in a cold water bath to room temperature and measure the absorbance at 540 nm against distilled water.

2. Enzyme blank

Incubate 1 ml of β -glucan substrate at +30°C for 10 minutes. Add 3 ml of DNS-solution and stir. Add 1 ml of enzyme dilution (in 0.1 M sodium acetate buffer, pH 5.0) and stir. Boil the mixture for exactly 5 minutes. Cool the reaction mixture in cold water bath to room temperature and measure the absorbance at 540 nm against distilled water.

The absorbance difference between the enzyme sample and the enzyme blank should be 0.3-0.5.

3. Standard curve

Prepare standard solutions from anhydrous glucose in distilled water. Glucose concentration in the standards should be 0.1-0.6 mg/ml. Pipette 1 ml of glucose standard solution into each of distilled water.

for exactly 5 minutes. Cool in a cold water bath to room temperature and measure the absorbance at 540 nm against standard blank. In the standard blank, glucose solution is replaced by 1 ml of distilled water. Otherwise standard blank is treated like glucose standard.

Plot glucose concentration as a function of absorbance. New standard curve is prepared for every new DNS-reagent.

Calculation

The β -glucanase activity of the sample is calculated according to the following equation:

$$\text{Activity (U/g)} = \frac{[A(X) - A(O)] \times k + C_0}{MW_{\text{glu}} \times t} \times 1000 \times Df$$

wherein:

A(X)	=	absorbance of the enzyme sample
A(O)	=	absorbance of the enzyme blank
k	=	the slope of the standard curve
C ₀	=	the intercept of glucose standard curve
1000	=	factor, mmol -> μ mol
Df	=	dilution factor (ml/g)

MW_{glu} = molecular weight of glucose
(180.16 mg/mmol)
 t = reaction time (10 minutes)

Based upon the above considerations, it is an object of the present invention to provide an enzyme feed additive for improving the FCR and/or increasing the digestibility of a cereal-based feed.

Accordingly, the present invention provides an enzyme feed additive comprising:

- (i) a xylanase;
- (ii) a protease; and optionally
- (iii) a β -glucanase

wherein the ratio of the units of xylanase activity per g of the feed additive to the units of β -glucanase activity per g of the feed additive is 1:0-0.25.

It has been found that the inclusion of the above enzyme feed additive in the diet of an animal enables the animal to digest the diet more efficiently. Thus, the addition of the additive to a feed increases the proportion of feed protein and energy which the animal can derive from the feed. This

It is also possible to utilise this aspect of the present invention by modifying a conventional feed by reducing its energy, and/or protein, and/or amino acid content whilst simultaneously maintaining the same nutritional levels of energy, protein, and amino acids available to the animal. This means that the amounts of costly energy and protein supplements conventionally included in an animal feed can be reduced as compared to conventional feeds. Energy supplements include fat. Protein supplements include fish-meal, meat-meal, soya-bean, rapeseed or canola. This results in a significant reduction in the cost per unit weight of the animal feed without decreasing its nutritional value. Alternatively, or even additionally, the amounts of amino acid supplements can be reduced as compared to conventional feeds which can also result in significant cost savings.

The enzyme feed additive according to the present invention can be prepared in a number of ways. For instance, it can be prepared simply by mixing different enzymes having the appropriate activities to produce an enzyme mix. This enzyme mix can be either mixed directly with a feed, or more conventionally impregnated onto a cereal-based carrier material such as milled wheat, maize or soya flour. Such an impregnated carrier also constitutes an enzyme feed additive in accordance with the present invention.

As an alternative, a cereal-based carrier formed from e.g. milled wheat or maize can be impregnated either simultaneously or sequentially with enzymes having the

appropriate activities. For example, a milled wheat carrier may be sprayed firstly with a xylanase, secondly with a protease, and optionally finally with a β -glucanase. The carrier material impregnated with these enzymes also constitutes an enzyme feed additive in accordance with the present invention.

The feed additive of the present invention may be mixed directly with the animal feed, or alternatively mixed with one or more other feed additives such as a vitamin feed additive, a mineral feed additive and an amino acid feed additive. The resulting feed additive including several different types of components can then be mixed in an appropriate amount with the feed.

The feed additive of the invention including the cereal-based carrier is normally mixed in amounts of 0.01-50 g per kilo of feed, more preferably 0.1-10 g/kilo and most preferably about 1 g/kilo.

An alternative way of preparing the enzyme feed additive of the present invention is to construct by recombinant DNA techniques a microorganism which produces the desired enzymes in the desired relative amounts. This can be done for instance by increasing the copy number of the gene encoding xylanase and/or by using a suitably strong promoter in front of the xylanase gene. Alternatively or additionally the microorganism strain can be deleted for certain cellulase genes especially endoglucanases.

The enzyme feed additive provided by the present invention may also include other enzymes such as one or more of an α -amylase, a glucoamylase, a pectinase, a mannanase, an α -galactosidase, a phytase and a lipase. Enzymes having the desired activities may for instance be mixed with the xylanase and protease either before impregnating these on a cereal-based carrier or alternatively such enzymes may be impregnated simultaneously or sequentially on such a cereal-based carrier. The carrier is then in turn mixed with a cereal-based feed to prepare the final feed. It is also possible to formulate the enzyme feed additive as a solution of the individual enzyme activities and then mix this solution with a feed material pre-formed as pellets or as a mash.

It is also possible to include the enzyme feed additive in the animal's diet by incorporating it into a second (and different) feed or drinking water which the animal also has access to. Accordingly, it is not essential that the enzyme mix provided by the present invention is incorporated into the cereal-based feed itself, although such incorporation forms a particularly preferred aspect of the present invention.

The enzyme mix provided by the present invention includes a xylanase as an essential component. This xylanase can be obtained from a bacterium such as *Bacillus*, *Streptomyces*, *Clostridium*, *Thermonospora*, *Microtetra-spora*, or *Ruminococcus*. However, it is more preferred that the xylanase is obtained from a fungus such as *Trichoderma*,

Aspergillus, *Humicola* or *Neocallimastix*. It is particularly preferred that the xylanase is the low pI xylanase (pI=5.2) and/or the high pI xylanase (pI=9.0) obtained from *Trichoderma longibrachiatum* obtainable by the method of Example 22 of WO 92/06209. It is particularly preferred that the xylanase is the high pI xylanase. Such a xylanase composition has a ratio of units of xylanase activity per unit amount to the units of β -glucanase activity per same unit amount of about 1:0.005. The xylanase may be a mutant xylanase having an amino acid sequence not found in nature, such a sequence corresponding to that of a naturally occurring xylanase modified by inserting, deleting or replacing one or more amino acid residues in the naturally occurring xylanase.

It is readily understood by those skilled in the art of enzyme production that enzyme compositions such as those described herein can be prepared by several different methods. For example, an enzyme composition including a xylanase and possessing only a limited amount of β -glucanase activity, can be produced by genetically engineering a host organism expressing such xylanase such that undesired genes are deleted or modified to inactivate the gene or the expression product therefrom. Additionally, the enzyme composition may be prepared by purification methods such as purifying a specifically desirable activity (xylanase) from a whole cellulase complex. Suitable methods are described in WO 92/06209. Other purification methods which are described in US Patent 5,328,841 include purification using a cellulase

be prepared by fermentation optimization procedures whereby the ratio of enzyme component activities can be modified by variation of the pH, temperature, carbon source, or combinations of these, during the fermentation procedure.

According to a most preferred aspect of the present invention, the xylanase is from *T. longibrachiatum* over-expressing high pI or low pI xylanase produced from a transformed host that is lacking one or more of functional endoglucanase I (EGI), functional endoglucanase II (EGII), functional cellobiohydrolase I (CBHI) and functional cellobiohydrolase II (CBHII). The preparation of such transformed hosts and the production of high and low pI xylanases is described in WO 92/06209 mentioned above.

The xylanase which results from such a host has significantly lower levels of enzymes possessing $\beta(1,4)$ -glucanase activity compared to xylanases obtained from naturally occurring sources. By using such a host, an enzyme composition can be obtained including only 0.005 units of β -glucanase activity per unit amount compared to 1 unit of xylanase activity in the same unit amount. In a preferred aspect of this invention, the enzyme feed additive includes a ratio of xylanase activity to β -glucanase activity of 1:0-0.01 and more preferably 1:0-0.005.

The ratio of the units of xylanase activity per g of the feed additive to the units of protease activity per g of the feed additive is preferably 1:0.001-1,000, more preferably 1:0.01-100 and most preferably 1:0.1-10.

The enzyme mix provided by the present invention also includes a protease as an essential component. It is preferred that the protease is a subtilisin which can be derived from the genus *Bacillus*, such as the strains including but not limited to *Bacillus amyloliquefaciens*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus subtilis*, or *Bacillus alcalophilus*.

Suitable proteases include but are not limited to the following commercially available proteases: Novo NEUTRASE (TM) (commercially available from Novo Nordisk); PURAFECT (TM) (commercially available from Genencor International, Inc); SAVINASE (TM) (commercially available from Novo Nordisk); MAXACAL (TM) (commercially available from Gist-Brocades); DURAZYM (TM) (commercially available from Novo Nordisk); and MAXAPEM (TM) (commercially available from Gist-Brocades).

The subtilisin may also be a mutant subtilisin having an amino acid sequence not found in nature, such a sequence corresponding to that of a naturally occurring subtilisin modified by inserting, deleting or replacing one or more amino acid residues in the naturally occurring subtilisin. Suitable mutant subtilisins are described in EP-A-0130756 (corresponding to US-Re-34606) (including mutations at position +155, +104, +222, +166, +33, +169, +189, +217, +156, +152); EP-A-0251446; WO91/06637 etc. The most preferred subtilisin is a mutant subtilisin having an amino acid sequence as described in EP-A-0130756.

tyr+217 of the subtilisin obtainable from *Bacillus amyloliquefaciens* with leucine.

Methods of producing such mutant subtilisins are described in detail in the publications US-Re-34606 and EP-A-0251446.

The protease which is included in the enzyme mix of the present invention should be free or relatively free of β -glucanase activity. This is to ensure that the resulting enzyme feed additive prepared by mixing the protease with the xylanase as previously described has the low or nil level of desired β -glucanase activity. This is less of a problem than with the xylanase component of the feed additive because microorganisms which naturally produce proteases do not necessarily produce high levels of β -glucanases.

It is found that the combination of the xylanase and the protease provided as the enzyme feed additive of the present invention have complementary and synergistic efficacies in terms of their ability to augment each other's effects to provide the advantages obtainable by the present invention of reducing the FCR of cereal-based animal feeds.

As mentioned above, the enzyme mix provided by the present invention is preferably for use as a feed additive in the preparation of a cereal-based feed.

According to a further aspect of the invention, this cereal-based feed comprises at least 25% by weight, more preferably at least 35% by weight, of wheat or maize or a combination of

both of these cereals. The feed further comprises a xylanase in such an amount that the feed includes 100-100,000 units of xylanase activity per kg.; a protease in such an amount that the feed includes 100-100,000 units of protease activity per kg.; and optionally a β -glucanase. The ratio of the units of xylanase activity per kg. of the feed to the units of β -glucanase activity per kg. of the feed falls within the range 1:0-0.25.

Preferably, the amount of feed additive added to the feed is such that the resulting feed comprises respectively 1,000-10,000 units of both xylanase activity and protease activity per kg.

In a further aspect of the invention, a cereal-based feed is provided which comprises at least 25% by weight of wheat and/or maize and 100-100,000 units per kg. of a mutant subtilisin comprising a substitution at the amino acid residue position equivalent to tyr+217 of the subtilisin obtainable from *Bacillus amyloliquefaciens* with leucine.

Cereal-based feeds according to the present invention are suitable for animals such as turkeys, geese, ducks, pigs, sheep and cows. The feeds though are particularly suitable for poultry and pigs, and in particular broiler chickens.

The present invention in a further aspect provides the use of a composition including the low pI xylanase and/or the high pI xylanase obtainable from *Trichoderma reesei*.

that the ratio of the units of xylanase activity per g of the composition to the units of β -glucanase activity per g of the composition is 1:0-0.25.

The above use of the composition as a feed additive produces a cereal-based feed having an improved FCR. According to a further aspect of the present invention, such a feed comprises (i) at least 25% by weight of wheat and/or maize; and (ii) the high pI xylanase and/or the low pI xylanase obtainable from *Trichoderma longibrachiatum* in such an amount that the feed includes 100-100,000 units of xylanase activity per kg.

The above use and cereal-based feed incorporating the high and/or low pI xylanase enables the animal to digest its feed more efficiently. Thus, the addition of such a xylanase containing feed additive to a feed increases the proportion of feed protein and energy which the animal can nutritionally derive from the feed. This in turn improves the FCR of the feed making it more economical in use. The feed additive including such a xylanase can of course be produced in the same way as the feed additives previously described.

The present invention is further explained by way of the following Examples.

Example 1

Cobb male broiler chickens were fed with the wheat-based starter feed set out in Table 1 below up to 21 days of age.

Table 1

Ingredients	Concentration (g/kg)
Wheat	637.1
Soybean meal	300.0
Limestone	13.3
Dicalcium Phosphate	13.0
DL-Methionine	3.1
Arginine	1.2
L-Lysine HCl	1.3
Vitamin mix	1.0
Mineral mix	1.0
Corn Oil	35.0
Salt	4.0

Calculated Composition

ME Kcal/kg	3004.0
CP %	22.8
Calcium %	0.87
Available Phosphorus	0.40
Methionine	0.64
Methionine + Cystine	1.01
Lysine	1.29

The above starter diet of the chickens was supplemented by varying amounts of xylanase obtained from *Trichoderma viride*

selected for relatively high xylanase activity such that it produces 55 units of β -glucanase activity to 1,000 units of xylanase activity per unit amount. The chickens were divided into 24 separate groups with 30 birds in each group. The diets of the 24 groups were supplemented by one of the combinations of 0, 500, 1,000, 2,000, 4,000 and 8,000 units/kg of xylanase; and 0, 250, 500 and 1,000 units/kg of β -glucanase.

The feed conversion ratio for each of these trials was obtained and the values of this are set out in the following Table 2:

Table 2

β -glucanase (units/kg) added	0	250	500	1,000	Average
Xylanase (units/kg)					
0	1.54	1.54	1.48	1.53	1.52
500	1.52	1.51	1.51	1.50	1.51
1000	1.47	1.49	1.52	1.54	1.51
2000	1.45	1.48	1.51	1.49	1.48
4000	1.50	1.51	1.46	1.49	1.49
8000	1.55	1.58	1.54	1.51	1.55

It is evident from the data set out in the above Table 2, that the presence of β -glucanase activity disadvantageously increases FCR values particularly, at the optimum xylanase level of around 1,000 - 3,000 units/kg.

From the results set out in the above Table 2, it is concluded that the units of xylanase activity present in the feed should be at least four times greater than the units of β -glucanase activity in order that an improvement in the FCR value can be obtained. Preferably the units of xylanase activity should be at least 100 times greater than the units of β -glucanase activity, and more preferably at least 1,000 times greater.

Example 2

Several groups of Ross 1 broiler chickens (mixed sex) were fed the starter and finisher diets set out in Table 3 below for days 0-21 and 22-42 of their life. Each of these groups included 90 birds.

Table 3

Ingredient (kg/t)	Starter	Finisher
Wheat	612.2	668.9
Soyabean meal 48% C.P.	318.8	240.9
Soya oil	32.6	55.4
Salt	3.03	3.01
DL methionine	2.01	0.4
Limestone	13.8	14.7
Dicalcium phosphate	12.5	11.7
Vitamin/mineral supplement	10.0	10.0

Calculated analysis(%)	Starter	Finisher
Crude protein	23.0	20.0
Calcium	0.90	0.90
Total phosphorus	0.67	0.63
Available phosphorus	0.42	0.40
Fat	4.63	6.82
Fibre	2.58	2.46
Lysine	1.21	0.99
Methionine	0.54	0.33
Sodium	0.15	0.15
Potassium	0.92	0.78
Chloride	0.24	0.24

A control group of broiler chickens did not have their diets supplemented by any enzymes. A second group Z had their diets supplemented by 3,000 units/kg of high pI xylanase produced from a transformed strain of *T. longibrachiatum* lacking functional EGI, EGII, CBHI and CBHII as described in WO 92/06209. Groups A, C, E had their diet supplemented with the same xylanase, and in addition by 2,000, 4,000 and 6,000 units/kg respectively of Novo Neutrase (TM) obtained from Novo Nordisk. Groups B, D and F had their diets supplemented with the same high pI xylanase, and in addition by 2,000, 4,000 and 6,000 units/kg respectively of a mutant subtilisin which is substituted at the amino acid residue position equivalent to tyr+217 of *Bacillus amyloliquefaciens* subtilin with leucine as described in the US-Re.-34606. The results of these tests are set out in the following Table 4.

Table 4

Group	Control group	Z	A	B	C	D	E	F
Final Body weight, kg	2.21	2.20	2.20	2.20	2.22	2.19	2.23	2.20
Feed Intake, g per bird per day	93.1	94.5	94.6	92.4	93.2	92.2	94.2	92.7
FCR (mortality adjusted)	1.85	1.85	1.85	1.82	1.85	1.82	1.82	1.82

From the above data, it can be calculated that the average FCR when using the xylanase and the protease Novo Neutrase is 1.84. On the other, the average FCR when using the xylanase and the mutant subtilisin protease of the present invention is 1.82. Each of the groups Z and A-F give rise to FCR values which are either equal to or superior to that obtained from the control group.

Example 3

Two groups of Ross 1 male broiler chickens were each fed the wheat-based starter feed set in Table 1 of Example 1 for days 0-7. The starter feed was not supplemented with any antibiotic, anticoccidial or any enzyme.

For days 7-21, each group was fed the basic wheat-based feed set out in the following Table 5 in the form of a mash:

Table 5

Ingredient	Amount (wt.%)
Soft wheat	65.5
Soybean meal 48% C.P.	27.3
Soy oil	3.1
Salt	0.3
DL methionine	0.2
Limestone	1.4
Dicalcium phosphate	1.2
Vitamin/mineral supplement	1.0

A first control group of the chickens did not have their diet supplemented by any enzymes. The above wheat-based feed fed to the second test group was supplemented with enriched low pI xylanase obtained from a *Trichoderma longibrachiatum* strain lacking functional EGI, EGII, CBHI and CBHII as described in WO 92/06209. The xylanase inclusion level in the wheat diet was 184 units/kg. The control group had an average FCR of 2.00. In contrast, the test group fed the feed supplemented with low pI xylanase had an FCR of 1.89. Such a reduction in the FCR of the feed by addition of the xylanase indicates that the nutritional performance of the feed is improved.

Example 4

Six groups each containing 12 castrated male pigs of PIC commercial genotype were respectively fed six different diets *ad libitum* in pellet form to assess the effect of enriched high pI xylanase obtained by *Trichoderma longibrachiatum*. At the start of the trial the pigs were all around 10 weeks of age. Thus, three basic feeds have a different maize/wheat ratio were formulated in accordance with the following Table 6 in which the amounts of the various components are expressed in terms kg/t of the feed.

Table 6

	20% Wheat diet	40% Wheat diet	60% Wheat diet
Maize	424	224	19
Wheat	200	400	600
Wheat Middlings	100	100	100
Soyabean meal (44)	200	200	200
Meat and bone meal	50	50	50
Canola oil	-	-	5
L-Lysine HCl	1.35	1.15	0.95
Vitamins & Minerals	25	25	25

Each diet was tested either as a control (without enzyme

high pI xylanase obtained from a *Trichoderma longibrachiatum* strain lacking functional EGI, EGII, CBHI and CBHII as described in WO 92/06209.

The results of these various tests are set in the following Table 7.

Table 7

Wheat(%)	20		40		60	
Maize(%)	42		22		2	
	Control feed	Feed plus xylanase	Control feed	Feed plus xylanase	Control feed	Feed plus xylanase
Start Weight (kg)	22.9	22.6	21.8	21.8	24.6	23.4
Finish Weight (kg)	75.0	82.7	78.7	82.2	78.8	82.8
Daily gain (g)	930	1073	1015	1076	974	1061
Daily feed intake (g)	2546	2773	2680	2638	2826	2942
FCR	2.76	2.59	2.68	2.47	2.93	2.81

It will be seen from the results set out in Table 7 that the addition of the xylanase significantly improved daily live weight gain by an average of 9% in finisher pigs offered diets containing 60% by weight of cereals formed from different proportions of maize and wheat. Overall, both feed intake and the FCR were significantly improved.

Example 5

Four groups of Nicholas male turkey poults were fed the maize-based starter feed described in Table 8 in the form of a mash up to 21 days of age.

Table 8

Ingredient	Amount (wt.%)
Maize	36.65
Soybean meal (45.6% CP)	55.4
Animal-vegetable fat	3.2
Dicalcium phosphate	2.3
Limestone	1.5
Mineral premix	0.3
Vitamin premix	0.3
Sodium chloride	0.15
DL methionine	0.2

The first control group were fed the feed of Table 8 unsupplemented. The feeds of three other test groups were supplemented respectively with 2,000 units/kg, 4,000 units/kg and 6,000 units/kg of a mutant subtilisin substituted at the amino residue position equivalent to tyr+217 of the *Bacillus amyloliquefaciens* subtilisin with leucine as described in US-Re.-34606. The results of these tests are set out in the following Table 9.

Table 9

Dietary Treatment	21-day Body Weight (grams/poult)	FCR
Control Feed	570	1.39
Feed + 2000 U/kg of protease	543	1.37
Feed + 4000 U/kg of protease	570	1.36
Feed + 6000 U/kg of protease	566	1.35

From the above data, it can be seen that increasing contents of the protease when added to the feed has the beneficial result of reducing the FCR of the feed.

The effect demonstrated above of improving FCR values can be obtained when feeds prepared in accordance with the present invention are fed to animals such as geese, ducks, sheep and cows, as well as to chickens, turkeys and pigs.

Claims:

1. An enzyme feed additive comprising:
 - (i) a xylanase;
 - (ii) a protease; and optionally
 - (iii) a β -glucanasewherein the ratio of the units of xylanase activity per g of the feed additive to the units of β -glucanase activity per g of the feed additive is 1:0-0.25.
2. An enzyme feed additive according to claim 1, wherein the xylanase is obtained from a bacterium.
3. An enzyme feed additive according to claim 2, wherein the bacterium is *Bacillus*, *Streptomyces*, *Clostridium* or *Ruminococcus*.
4. An enzyme feed additive according to claim 1, wherein the xylanase is obtained from a fungus.
5. An enzyme feed additive according to claim 4, wherein the fungus is *Trichoderma*, *Aspergillus*, *Humicola* or *Neocallimastix*.
6. An enzyme feed additive according to claim 5, wherein the xylanase is the low pI xylanase and/or the high pI xylanase.

7. An enzyme feed additive according to claim 6, which comprises the high pI xylanase.
8. An enzyme feed additive according to claim 1, wherein the xylanase is a mutant xylanase having an amino acid sequence not found in nature, such a sequence corresponding to that of a naturally occurring xylanase modified by inserting, deleting or replacing one or more amino acid residues in the naturally occurring xylanase.
9. An enzyme feed additive according to any preceding claim, wherein the protease is a subtilisin.
10. An enzyme feed additive according to claim 9, wherein the subtilisin is derived from the genus *Bacillus*.
11. An enzyme feed additive according to claim 10, wherein the subtilisin is derived from *Bacillus amyloliquefaciens*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus subtilis*, or *Bacillus alcalophilus*.
12. An enzyme feed additive according to claim 9, wherein the subtilisin is a mutant subtilisin having an amino acid sequence not found in nature, such a sequence corresponding to that of a naturally occurring subtilisin modified by inserting, deleting or replacing one or more amino acid residues in the naturally occurring subtilisin.

13. An enzyme feed additive according to claim 12, wherein the mutant subtilisin comprises a substitution at the amino acid residue position equivalent to asn+155, tyr+104, met+222, gly+166, ser+33, gly+169, phe+189, tyr+217, glu+156, or ala+152 of *Bacillus amyloliquefaciens* subtilisin with one of the other 19 naturally occurring amino acids.
14. An enzyme feed additive according to claim 13, wherein the mutant subtilisin comprises a substitution at the amino acid residue position equivalent to tyr+217 of *Bacillus amyloliquefaciens* subtilisin with leucine.
15. An enzyme feed additive according to any preceding claim, wherein the ratio of the units of xylanase activity to β -glucanase activity is 1:0-0.01.
16. An enzyme feed additive according to claim 15, wherein the ratio of the units of xylanase activity to β -glucanase activity is 1:0-0.005.
17. An enzyme feed additive according to any preceding claim, wherein the ratio of the units of xylanase activity per g of the feed additive to the units of protease activity per g of the feed additive is 1:0.001 - 1,000
18. An enzyme feed additive according to any preceding claim, wherein the ratio of the units of xylanase activity per g of the feed additive to the units of protease activity per g of the feed additive is 1:0.001 - 1,000

glucoamylase, pectinase, mannanase, α -galactosidase, phytase and lipase.

19. An enzyme feed additive according to any preceding claim, further comprising a carrier.
20. An enzyme feed additive according to claim 19, wherein the carrier is milled wheat, maize, soya or a by-product of any of these materials.
21. The use of an enzyme feed additive according to any preceding claim, for improving the feed conversion ratio and/or increasing the digestibility of a cereal-based feed.
22. A cereal-based feed comprising (i) at least 25% by weight of wheat and/or maize; (ii) a xylanase in such an amount that the feed includes 100-100,000 units of xylanase activity per kg.; (iii) a protease in such an amount that the feed includes 100-100,000 units of protease activity per kg.; and optionally (iv) a β -glucanase; wherein the ratio of the units of xylanase activity per kg. of the feed to the units of β -glucanase activity per kg. of the feed is 1:0-0.25.
23. A cereal-based feed according to claim 22, which comprises 1,000-10,000 units of xylanase activity per kg. and 1,000-10,000 units of protease activity per kg.

24. A cereal-based feed according to claim 22 or claim 23, comprising at least 35% by weight of wheat and/or maize.
25. A cereal-based feed comprising at least 25% by weight of wheat and/or maize and 100-100,000 units per kg. of a mutant subtilisin comprising a substitution at the amino acid residue position equivalent to tyr+217 of the subtilisin obtainable from *Bacillus amyloliquefaciens* with leucine.
26. A cereal-based feed comprising (i) at least 25% by weight of wheat and/or maize; and (ii) the high pI xylanase and/or the low pI xylanase obtainable from *Trichoderma longibrachiatum* in such an amount that the feed includes 100-100,000 units of xylanase activity per kg.
27. Use of a cereal-based feed according to any of claims 22-26 as a poultry feed or as a pig feed.
28. Use of a composition including the low pI xylanase and/or the high pI xylanase obtainable from *Trichoderma longibrachiatum* and optionally a β -glucanase as a feed additive, characterised in that the ratio of the units of xylanase activity per g of the composition to the units of β -glucanase activity per g of the composition is 1:0-0.25.

INTERNATIONAL SEARCH REPORT

Int. Application No.
PCT/EP 95/03277

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 A23K1/165 A23K1/18		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 6 A23K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATENT ABSTRACTS OF JAPAN vol. 9 no. 205 (C-299) ,22 August 1985 & JP,A,60 075238 (TOSHIO TSUBAKI) 27 April 1985, cited in the application see abstract ---	1,22
A	GB,A,2 261 877 (KYOWA HAKKO KOGYO CO. LTD.) 2 June 1993 see page 3, line 1 - page 4, line 3; claims 1,2 ---	1
A	WO,A,92 06209 (GENENCOR INTERNATIONAL INC.) 16 April 1992 cited in the application see figure 16; example 22 ---	1,4-8
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 30 November 1995		Date of mailing of the international search report 20.12.95
Name and mailing address of the ISA European Patent Office P.B. 581b Patenlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340.2040, Tx. 31 651 epo nl, Fax: (+ 31-70) 340-1016		Authorized officer Alvarez Alvarez, C

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Int. Application No.
PCT/EP 95/03277

C (Continuation): DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 463 706 (GIST-BROCADES N.V.) 2 January 1992 see claims 1-3,16; example 8 ---	4,5,19, 22
A	EP,A,0 507 723 (NOVO NORDISK A/S) 7 October 1992 see page 3, line 39 - line 52 ---	1,4,5
A	WO,A,93 25693 (G. HAZLEWOOD ET AL.) 23 December 1993 see abstract ---	4,5
A	WO,A,91 04673 (NOVO NORDISK A/S ET AL.) 18 April 1991 cited in the application see example 6 ---	4-7,20, 21,24, 26,27
A	WO,A,92 10945 (ENZYMATIX LTD. ET AL.) 9 July 1992 see abstract ---	1
A	EP,A,0 130 756 (GENENTECH INC.) 9 January 1985 cited in the application see page 1, line 20 - page 2, line 22 ---	1,9-14
A	EP,A,0 251 446 (GENENTECH INC.) 7 January 1988 cited in the application see abstract ---	1,9-14
A	WO,A,91 06637 (GENENCOR INTERNATIONAL) 16 May 1991 cited in the application see abstract -----	1,9-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 95/03277

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
GB-A-2261877	02-06-93	DE-A-	4239139	27-05-93
		JP-A-	5192093	03-08-93

WO-A-9206209	16-04-92	CA-A-	2093421	06-04-92
		CA-A-	2093424	06-04-92
		CA-A-	2093428	06-04-92
		EP-A-	0551386	21-07-93
		EP-A-	0551394	21-07-93
		EP-A-	0552276	28-07-93
		EP-A-	0553280	04-08-93
		FI-A-	931490	31-05-93
		FI-A-	931491	25-05-93
		FI-A-	931495	27-05-93
		JP-T-	6502223	10-03-94
		JP-T-	6502205	10-03-94
		JP-T-	6511379	22-12-94
		JP-T-	6502226	10-03-94
		US-A-	5419778	30-05-95
		WO-A-	9206210	16-04-92
		WO-A-	9206221	16-04-92
		WO-A-	9206183	16-04-92
		US-A-	5246853	21-09-93
		US-A-	5290474	01-03-94
		US-A-	5328841	12-07-94

EP-A-463706	02-01-92	AT-T-	124844	15-07-95
		AU-B-	654147	27-10-94
		AU-B-	8318691	18-02-92
		AU-B-	647170	17-03-94
		AU-B-	8320591	18-02-92
		CA-A-	2067329	25-01-92
		DE-D-	69111148	17-08-95
		EP-A-	0468596	29-01-92
		JP-T-	5501657	02-04-93
		JP-T-	5500907	25-02-93
		WO-A-	9201389	06-02-92
		WO-A-	9201793	06-02-92
		US-A-	5358864	25-10-94

EP-A-507723	07-10-92	WO-A-	9217573	15-10-92

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 95/03277

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-507723		EP-A- 0579672	26-01-94
		JP-T- 6506348	21-07-94
WO-A-9325693	23-12-93	AU-B- 4298393	04-01-94
		AU-B- 4347993	04-01-94
		WO-A- 9325671	23-12-93
		CA-A- 2138383	23-12-93
		CA-A- 2138399	23-12-93
		EP-A- 0652961	17-05-95
		FI-A- 945889	08-02-95
		FI-A- 945928	16-02-95
		JP-T- 7508647	28-09-95
		NO-A- 944851	14-02-95
WO-A-9104673	18-04-91	DE-D- 69007115	07-04-94
		DE-T- 69007115	09-06-94
		EP-A 0494916	22-07-92
		JP-T- 5500807	18-02-93
WO-A-9210945	09-07-92	AT-T- 126972	15-09-95
		AU-B- 9062291	22-07-92
		DE-D- 69112643	05-10-95
		EP-A- 0563133	06-10-93
		US-A- 5432074	11-07-95
EP-A-130756	09-01-85	US-A- 4760025	26-07-88
		AU-B- 587960	07-09-89
		AU-B- 2956884	03-01-85
		AU-B- 631797	10-12-92
		AU-B- 3714989	23-11-89
		AU-B- 636109	22-04-93
		AU-B- 3720889	07-12-89
		DE-A- 3486139	03-06-93
		DE-T- 3486139	11-11-93
		EP-A, B 0247647	02-12-87
		EP-A, B 0246678	25-11-87
		EP-A- 0357157	07-03-90
		JP-A- 60070075	20-04-85
		JP-A- 6319534	22-11-94
		JP-A- 6315378	15-11-94

INTERNATIONAL SEARCH REPORT

information on patent family members

Int. Application No
PCT/EP 95/03277

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-130756		NZ-A- 236052	25-03-94
		NZ-A- 244586	28-08-95
		US-A- 5310675	10-05-94
		US-A- 5371008	06-12-94
		US-A- 5371190	06-12-94
		US-A- 5185258	09-02-93
		US-A- 5182204	26-01-93
		US-A- 5352594	04-10-94
		US-A- 5204015	20-04-93
		US-A- 5346823	13-09-94
		US-A- 5441882	15-08-95
		US-E- RE34606	10-05-94
		US-A- 5244791	14-09-93
		US-A- 5411873	02-05-95
		US-A- 5264366	23-11-93
EP-A-251446	07-01-88	AT-T- 116365	15-01-95
		AU-B- 614929	19-09-91
		AU-B- 7228187	05-11-87
		DE-D- 3750916	09-02-95
		DE-T- 3750916	01-06-95
		ES-T- 2068181	16-04-95
		JP-A- 1085075	30-03-89
		NZ-A- 240670	26-05-95
		US-A- 5371008	06-12-94
		US-A- 5371190	06-12-94
		US-A- 5185258	09-02-93
		US-A- 5182204	26-01-93
		US-A- 5204015	20-04-93
		US-A- 5352594	04-10-94
WO-A-9106637	16-05-91	US-A- 5185258	09-02-93
		AU-B- 633376	28-01-93
		AU-B- 6634190	31-05-91
		CA-A- 2044630	01-05-91
		CN-A- 1052897	10-07-91
		EP-A- 0451244	16-10-91
		IE-B- 62771	22-02-95
		JP-T- 4505263	17-09-92
		US-A- 5204015	20-04-93

information on patent family members

PCT/EP 95/03277

Form FD-35A (10-6-65) (Instructions on back) (Rev. 1-25-64)

